Magnosphere™ MS300/Streptavidin

PRODUCT DESCRIPTION

Magnosphere™ MS300/Streptavidin beads are magnetic microparticles coated with streptavidin for bioseparation. The particle surfaces are covered with a JSR Life Sciences proprietary hydrophilic polymer to give the beads their characteristic low non-specific binding without inhibition of enzyme activities. Consequently, *Magnosphere™ MS300/Streptavidin* beads can be used for a variety of applications such as PCR or immunoassays for excellent performance.

FEATURES

- High affinity to biotinylated proteins and nucleotides
- Uniform particle size
- Superparamagnetic
- Rapid magnetic responsiveness
- Low non-specific binding

EXAMPLE APPLICATIONS

PCR, quantitative PCR, Immunoassay Immobilization of biotinylated DNA/RNA, Immobilization of biotinylated proteins

SPECIFICATIONS

Package volume	2 mL
Solid content in slurry	1 % (10 mg/mL, 6 x 10 ⁸ beads/mL approx.)
Dispersion media	TBS* + 0.05 % Tween20 + 0.09 % Sodium Azide
Bead diameter	3 μm (micrometer)
Bead magnetite content	20 % approx.
Biotin binding capacity	400-600 pmol/mg bead
Shelf life	Labeled on the bottle
*TBS: Tris buffered saline, 25	mM Tris-HCI (pH 7.2) / 0.15 M NaCl

STORAGE

Stored at 2-8 °C. Do not freeze the vial. Vortex the vial or pipette gently up and down to obtain a homogeneous dispersion before use.

DISPOSAL

Reagent contains sodium azide at a low concentration as a preservative. Sodium azide is toxic if ingested and may react with heavy metals to form explosive metal azides. Azide compounds should be diluted with running water before discarding to avoid deposits in plumbing where explosive condition may develop.

RECOMMENDED PROTOCOLS

[Protocol I] Immobilization of biotinylated DNA on *Magnosphere*™ *MS300/Streptavidin* beads.

Reagent and equipment requirement

Binding Buffer (2X): Equipment: 20 mM Tris-HCI (pH 7.4) with 1 mM EDTA, 2 M NaCl, 0.1 % Tween20 Magnetic separator. Vortex tube mixer. Tube rotator.

- 1. Suspend the *Magnosphere™ MS300/Streptavidin* well using Vortex mixer and put 100 μL of the suspension (i.e., 1 mg beads) into a microtube.
- 2. Place the tube on a magnetic tube stand for 1 minute (or longer if needed) and remove the supernatant carefully.
- 3. Add 200 μ L of 1x Binding Buffer and suspend the beads by vortexing. Then, remove the supernatant as in step 2.
- 4. Add biotinylated DNA solution [e.g. 5 µg of DNA] and an equal volume of 2x Binding Buffer to the microtube. Suspend the beads by vortexing.
- 5. Keep rotating the tube with Tube rotator for 10 minutes at room temperature.
- 6. Remove the supernatant as in step 2.
- 7. Wash the beads using 200 μ L of 1x Binding Buffer and suspend the beads by vortexing.
- 8. Remove the supernatant as in step 2.
- 9. Repeat steps 7 & 8 for a total of 3 times.
- 10. Suspend the beads with a desired buffer suitable for downstream applications and store at 2-8 °C until needed

EXPERIMENTAL EXAMPLE

[Example I] Influence of Magnosphere™ MS300/Streptavidin on qPCR

Various amounts of *Magnosphere™ MS300/Streptavidin* beads were added to a PCR cocktail containing DNA (100bp in length). After qPCR reaction, the Threshold Cycle (Ct) was plotted against



the amount of beads added to the reaction.

Conclusion: Addition of *Magnosphere*™ *MS300/Streptavidin* beads up to 3mg/tube does not affect the PCR reaction.

[Example II] Recovery of biotinylated dsDNA onto *Magnosphere*™ *MS300/Streptavidin*

Various amounts of biotinylated 100bp dsDNA were mixed with 1mg of *Magnosphere*™ *MS300/Streptavidin* beads in the Binding Buffer. The amounts of immobilized dsDNA were



quantitated through qPCR.

Conclusion: Added dsDNA was quantitatively recovered on the *Magnosphere*™ *MS300/Streptavidin* beads up to 20pmol (i.e., 10¹³ copies).



IMPORTANT NOTICE

- This product is for research use only and not intended for therapeutic or in vivo diagnostic use.
- The specifications of the product may be changed without a notice.
- JSR Life Sciences Corporation does not guarantee that this product will be continuously available.
- JSR Life Sciences Corporation makes no warranties as to this product including, but not limited to, implied warranties of merchantability or fitness for a particular purpose.

CONTACT INFORMATION

JSR Corporation Life Sciences Division E-mail: dx@jls.jsr.co.jp URL: http://www.jsrlifesciences.com/