

For Research Use Only.
Not for use in diagnostic procedures.

Anti-DDDDK-tag mAb-Alexa Fluor[®] 647

| | |
|--------------------|---|
| CODE No. | M185-A64 |
| CLONALITY | Monoclonal |
| CLONE | FLA-1 |
| ISOTYPE | Mouse IgG2a κ |
| QUANTITY | 100 μ L, 1 mg/mL |
| SOURCE | Purified IgG from hybridoma supernatant |
| IMMUNOGEN | KLH conjugated DYKDDDDK peptide |
| REACTIVITY | This antibody reacts with N-terminal, Internal and C-terminal DDDDK-tagged (DYKDDDDK) proteins. |
| FORMULATION | PBS containing 1% BSA and 0.1% ProClin 150. |
| STORAGE | This antibody solution is stable for one year from the date of purchase when stored at 4°C. |

APPLICATIONS-CONFIRMED

| | |
|----------------------------|------------------|
| <u>Immunocytochemistry</u> | 0.5-1 μ g/mL |
| <u>Flow cytometry</u> | 0.5 μ g/mL. |

RELATED PRODUCTS

Please visit our web site <https://ruo.mbl.co.jp/>.

LABEL LICENSES:


This product is provided under an agreement between LIFE TECHNOLOGIES Corporation, and Medical & Biological Laboratories Co., LTD.

Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc

The descriptions of the following protocols are examples. Each user should determine the appropriate condition.

Immunocytochemistry

- 1) Spread the cells in the nutrient condition on a glass slide, then incubate in a CO₂ incubator for one night.
- 2) Remove the culture supernatant by careful aspiration.
- 3) Fix the cells by immersing the slide in 4% paraformaldehyde (PFA)/PBS for 10 min. at room temperature (20~25°C).
- 4) Prepare a wash container such as a 500 mL beaker with a magnetic stirrer. Then wash the fixed cells on the glass slide by soaking the slide with a plenty of PBS in the wash container for 5 min. Take care not to touch the cells. Repeat another wash once more.
- 5) Immerse the slide in 0.2% Triton X-100/PBS for 10 min. at room temperature.
- 6) Wash the slide in a plenty of PBS as in the step 4).
- 7) Cover each cell with Clear Back (human Fc receptor blocking reagent, MBL; code no. MTG-001) for 5 min. at room temperature.
- 8) Add 200 µL of the primary antibody diluted with PBS as suggested in the **APPLICATIONS** onto the cells and incubate for 1 hr. at room temperature. (Optimization of antibody concentration or incubation condition is recommended if necessary.)
- 9) Wash the slide in a plenty of PBS as in the step 4).
- 10) Wipe excess liquid from slide but take care not to touch the cells. Never leave the cells to dry.
- 11) Counter stain with DAPI for 5 min. at room temperature.
- 12) Wash the slide in a plenty of PBS as in the step 4).
- 13) Promptly add mounting medium onto the slide, then put a cover slip on it.




Immunocytochemical detection of DDDDK-tagged protein in HeLa

Magenta: M185-A64

Cyan: DAPI

Flow cytometric analysis for adherent cells

- 1) Detach the cells from culture dish.
- 2) Wash the cells once with 1 mL of washing buffer [PBS containing 2% fetal calf serum (FCS)].
- 3) Add 200 μ L of 4% paraformaldehyde (PFA) to the cell pellet after tapping. Mix well, then fix the cells for 10 min. at room temperature.
- 4) Wash the cells twice with 1 mL of washing buffer.
- 5) Add 200 μ L of PBS containing 0.2% Triton X-100 to the cell pellet after tapping. Mix well, then permeabilize the cells for 10 min. at room temperature.
- 6) Wash the cells once with 1 mL of washing buffer.
- 7) Resuspend the cells with washing buffer (1.6×10^6 cells/mL).
- 8) Add 100 μ L of the cell suspension into each tube, and centrifuge at 500 x g for 1 min. at room temperature (20~25°C). Remove supernatant by careful aspiration.
- 9) Add 20 μ L of Clear Back (human Fc receptor blocking reagent, MBL; code no. MTG-001) to the cell pellet after tapping. Mix well and incubate for 5 min. at room temperature.
- 10) Add 40 μ L of the primary antibody at the concentration as suggested in the **APPLICATIONS** diluted in the washing buffer. Mix well and incubate for 30 min. at room temperature.
- 11) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 min. at room temperature. Remove supernatant by careful aspiration. Repeat another wash once more.
- 12) Resuspend the cells with 500 μ L of the washing buffer and analyze by a flow cytometer.



Flow cytometric detection of DDDDK-tagged protein in HeLa

Closed: M185-A64

Open: Isotype control (M076-A64)

Upper: DDDDK-tagged protein in HeLa

Lower: Parental cell (HeLa)