

Anti-Mincle (Mouse) mAb

CODE No.	D266-3
CLONALITY	Monoclonal
CLONE	1B6
ISOTYPE	Rat IgG1 κ
QUANTITY	100 μ L, 1 mg/mL
SOURCE	Purified IgG from hybridoma supernatant
IMMUNOGEN	RBL-2H3 cells expressing full-length mouse mincle
REACTIVITY	This clone reacts with mouse Mincle (Clec4e) and crossreacts weakly with MCL (Clec4d).
FORMURATION	PBS containing 50% Glycerol (pH 7.2). No preservative is contained.
STORAGE	This antibody solution is stable for one year from the date of purchase when stored at -20°C.

APPLICATIONS-CONFIRMED

<u>Western blotting</u>	1 μ g/mL for chemiluminescence detection system
<u>Immunoprecipitation</u>	2 μ g/2 x 10 ⁶ cells/sample
<u>Flow cytometry</u>	2 μ g/mL

APPLICATION-REPORTED

<u>Functional activity</u>	1-10 μ g/mL for blocking, Reference 1) and 2)
----------------------------	---

SPECIES CROSS REACTIVITY on WB

Species	Human	Mouse	Rat	Guinea Pig
Cell	Not tested	LPS-stimulated Balb/c mouse peritoneal macrophage	Not tested	Not tested
Reactivity		+		

Entrez Gene ID 56619 (Mouse)

- REFERENCES**
- 1) Yamasaki, S., *et al.*, *Nat. Immunol.* **9**, 1179-1188 (2008) [WB, FCM, Function]
 - 2) Miyake, Y., *et al.*, *Immunity* **38**, 1050-1062 (2013) [FCM, Function]
 - 3) Yamasaki, S., *et al.*, *PNAS* **106**, 1897-1902 (2009)
 - 4) Ishikawa, E., *et al.*, *J. Exp. Med.* **206**, 2879-2888 (2009)

For more information, please visit our web site <http://ruo.mbl.co.jp/>

RELATED PRODUCTS

D266-3	Anti-Mincle (Mouse) mAb (1B6)
D292-3	Anti-Mincle (Mouse) mAb (4A9)
D316-3	Anti-Mincle (Guinea pig) mAb (5H4)
M191-3	Anti-FcεR1γ (FcRγ) (Mouse) mAb (1D6)
M191-A48	Anti-FcεR1γ (FcRγ) (Mouse) mAb -Alexa Fluor® 488 (1D6)
M191-A59	Anti-FcεR1γ (FcRγ) (Mouse) mAb -Alexa Fluor® 594 (1D6)
M191-A64	Anti-FcεR1γ (FcRγ) (Mouse) mAb -Alexa Fluor® 647 (1D6)
PM068	Anti-FcεR1γ (FcRγ) (Mouse) pAb
D222-3	Anti-GITR (Mouse) mAb (DTA-1)
D203-3	Anti-IL-15 (Mouse) mAb (AIO.3)
D043-3	Anti-IL-18 (Human) mAb (25-2G)
D044-3	Anti-IL-18 (Human) mAb (125-2H)
D045-3	Anti-IL-18 (Human) mAb (159-12B)
D046-3	Anti-IL-18 (Mouse) mAb (39-3F)
D047-3	Anti-IL-18 (Mouse) mAb (74)
D048-3	Anti-IL-18 (Mouse) mAb (93-10C)
M138-3	Anti-IL-33 (Human) mAb (5H1)
M161-3	Anti-IL-33 (Mouse) mAb (4G4)
PM033	Anti-IL-33 (Human) pAb
D205-3	Anti-TLR4 (CD284) (Mouse) mAb (UT49)
D205-4	Anti-TLR4 (CD284) (Mouse) mAb-FITC (UT49)
D079-3	Anti-TLR4-MD-2 complex (Mouse) mAb (MTS510)
D206-3	Anti-TLR4-MD-2 complex (Mouse) mAb (UT15)
D077-3	Anti-TLR4 (CD284) (Human) mAb (HTA125)


Other related antibodies and kits are also available.

Please visit our website at <http://ruo.mbl.co.jp/>

SDS-PAGE & Western blotting

- 1) Wash 1×10^7 cells 3 times with PBS and suspend them in 1 mL of Laemmli's sample buffer, then sonicate briefly (up to 10 sec.).
- 2) Boil the samples for 2 min. and centrifuge. Load 10 μ L of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel (12.5% acrylamide) for electrophoresis.
- 3) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm² for 1 hr. in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 4) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) for 1 hr. at room temperature.
- 5) Incubate the membrane with primary antibody diluted with 1% skimmed milk (in PBS, pH 7.2) as suggested in the **APPLICATIONS** for 1 hr. at room temperature. (The concentration of antibody will depend on the conditions.)
- 6) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 min. x 3 times).
- 7) Incubate the membrane with the 1:10,000 of anti-IgG (Rat) pAb-HRP (MBL; code no. IM-0825) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hr. at room temperature.
- 8) Wash the membrane with PBS-T (5 min. x 3 times).
- 9) Wipe excess buffer on the membrane, and then incubate it with appropriate chemiluminescence reagent for 1 min. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 10) Expose to an X-ray film in a dark room for 3 min. Develop the film as usual. The condition for exposure and development may vary.

(Positive control for Western blotting; LPS-stimulated Balb/c mouse peritoneal macrophage)



Western blot analysis of mouse Mincle expressed in Balb/c mouse peritoneal macrophages

Lane 1: Non-stimulation
Lane 2: LPS-stimulated

Immunoblotted with Anti-Mincle (Mouse) mAb (D266-3)

Immunoprecipitation


[Preparation of antibody-bound protein G beads]

- a) Mix 20 μ L of 50% protein G agarose beads slurry resuspended in 500 μ L of IP buffer (10 mM Tris-HCl pH 8.0, 500 mM NaCl, 0.1% NP-40) with primary antibody as suggested in the **APPLICATIONS**. Incubate with gentle agitation for 1 hr. at room temperature.
- b) Wash the beads 3 times with 1 mL of IP Buffer.
- c) Wash the beads 3 times with 1 mL of Fixation Buffer (0.1 M $\text{Na}_2\text{B}_4\text{O}_7$)
- d) Add 200 μ L of 20 mM Dimethyl pimelimidate (DMP) in Fixation buffer. Incubate with gentle agitation for 30 min. at room temperature.
- e) Centrifuge the tube at 300 x g for 2 min. at 4°C. Carefully discard the sup and add 200 μ L of 200 mM Tris-HCl, pH 8.0. Incubate at 4°C until just before use.
- f) Wash the beads 3 times with 1 mL of 200 mM Tris-HCl, pH 8.0.
- g) Wash the beads 1 time with 1 mL of IP buffer.

[Protocol]

- 1) Wash 4 x 10⁶ cells 2 times with PBS and resuspend them with 1 mL of Extraction buffer (50 mM Tris-HCl pH 7.4, 150mM NaCl, 0.05% NP-40) containing appropriate protease inhibitors, then sonicate briefly (up to 15 sec.).
- 2) Incubate the tube for 10 min. on ice.
- 3) Centrifuge the tube at 12,000 x g for 5 min. at 4°C and transfer the supernatant to another tube.
- 4) Add 500 μ L of the supernatant to the tube containing antibody conjugated beads (Step g)).
- 5) Incubate with gentle agitation for 1 hr. at room temperature.
- 6) Wash the beads 6 times with 1 mL of Extraction buffer.
- 7) Resuspend the beads in 20 μ L of Laemmli's sample buffer, boil for 3 min. and centrifuge. Use 10 μ L/lane for the SDS-PAGE analysis. (See **SDS-PAGE & Western blotting**.)

(Positive control for Immunoprecipitation; LPS-stimulated Balb/c mouse peritoneal macrophage)



Immunoprecipitation of mouse Mincle from LPS-stimulated Balb/c mouse peritoneal macrophages

Lane 1: Isotype control (M080-3)


Lane 2: Anti-Mincle (Mouse) mAb (D266-3)

Immunoblotted with D266-3

Flow cytometric analysis

- 1) Wash the cells (2×10^5 cells/sample) 2 times with 1 mL of washing buffer (PBS containing 2% fetal calf serum (FCS)).
- 2) Add 20 μ L of 1 mg/mL human IgG in normal goat serum to the cell pellet after tapping. Mix well and incubate for 5 min. at room temperature.
- 3) Add 40 μ L of the primary antibody at the concentration as suggested in the **APPLICATIONS** diluted in the washing buffer. Mix well and incubate for 30 min. at room temperature.
- 4) Wash the cells 1 time with 1 mL of the washing buffer.
- 5) Add 40 μ L of 1:100 anti-IgG (Rat) pAb-FITC (MBL; code no. IM-0827) diluted with the washing buffer. Mix well and incubate for 15 min. at room temperature.
- 6) Wash the cells 1 time with 1 mL of the washing buffer.
- 7) Add 40 μ L of 1:100 CD11b (Mac-1)-PE (Beckman Coulter; code no. 732048) diluted with the washing buffer. Mix well and incubate for 15 min. at room temperature.
- 8) Wash the cells 1 time with 1 mL of the washing buffer.
- 9) Resuspend the cells with 500 μ L of the washing buffer and analyze by a flow cytometer.

(Positive control for Flow cytometry; LPS-stimulated Balb/c mouse peritoneal macrophage)



Flow cytometric detection of mouse Mincle in LPS-stimulated Balb/c mouse peritoneal macrophages

Left: Isotype control (M080-3)
Right: Anti-Mincle (Mouse) mAb (D266-3)