

# Anti-GP2 (Glycoprotein 2) (Mouse) mAb -Alexa Fluor<sup>®</sup> 488

<b>CODE No.</b>	D278-A48
<b>CLONALITY</b>	Monoclonal
<b>CLONE</b>	2F11-C3
<b>ISOTYPE</b>	Rat IgG2a $\kappa$
<b>QUANTITY</b>	100 $\mu$ L, 1 mg/mL
<b>SOURCE</b>	Purified IgG from hybridoma supernatant
<b>IMMUNOGEN</b>	Mouse GP2, extracellular domain (recombinant, human Fc fusion protein)
<b>FORMULATION</b>	PBS containing 1% BSA and 0.1% ProClin 150
<b>STORAGE</b>	This antibody solution is stable for one year from the date of purchase when stored at 4°C.

## APPLICATIONS-CONFIRMED

<u>Immunohistochemistry</u>	5 $\mu$ g/mL
<u>Flow cytometry</u>	1 $\mu$ g/mL

## SPECIES CROSS REACTIVITY

Species	Human	Mouse	Rat	Hamster
Tissues	Peyer's patches	Peyer's patches	Not tested	Not tested
Reactivity	-	+		

**Entrez Gene ID** 67133 (Mouse)

## REFERENCES

- 1) Laphorne, S., *et al.*, *Immunology*, in press
- 2) Donaldson, D. S., *et al.*, *Mucosal Immunol.* **5**, 216-225 (2012)
- 3) Fukuda, S., *et al.*, *J. Vis. Exp.* **58**, e3225 (2011)
- 4) Ebisawa, M., *et al.*, *Int. Immunol.* **23**, 261-269 (2011)
- 5) Hase, K., *et al.*, *Nature* **462**, 226-230 (2009)

This clone is used in these references.

For more information, please visit our web site <https://ruo.mbl.co.jp/>

## LABEL LICENSES:

This product is provided under an agreement between LIFE TECHNOLOGIES Corporation, and Medical & Biological Laboratories Co.,

LTD. Alexa Fluor<sup>®</sup> is a registered trademark of Molecular Probes, Inc.

## RELATED PRODUCTS


Please visit our web site <https://ruo.mbl.co.jp/>

The descriptions of the following protocols are examples. Each user should determine the appropriate condition.

### **Immunohistochemical detection for paraffin embedded section**

- 1) Deparaffinize the sections with Xylene 3 times for 3 min. each.
- 2) Wash the slides with Ethanol 3 times for 3 min. each.
- 3) Wash the slides 1 time in PBS-T (0.05% Tween-20 in PBS) for 5 min.
- 4) Remove the slides from PBS-T, wipe gently around each section and cover tissues with 0.5% blocking reagent (Parkin Elmer) in PBS for 30 min. to block non-specific staining. Do not wash.
- 5) Tip off the blocking buffer, wipe gently around each section and cover tissues with primary antibody diluted with 0.5% blocking reagent in PBS as suggested in the **APPLICATIONS**. (Optimization of antibody concentration or incubation condition is recommended if necessary.)
- 6) Incubate the sections overnight at 4°C.
- 7) Wash the slides 3 times in PBS-T for 5 min. each.
- 8) Now ready for mounting.

(Positive control for Immunohistochemistry; Mouse Peyer's patches)



### ***Immunohistochemical detection of mouse GP2 in Peyer's patches***


Strain: C57BL/6

Green: D278-A48

### **Flow cytometric analysis**

- 1) Wash the cells ( $5 \times 10^5$  cells/sample) 1 time with 1 mL of washing buffer [PBS containing 2% fetal calf serum (FCS)].
- 2) Add 20  $\mu$ L of Clear Back (human Fc receptor blocking reagent, MBL; code no. MTG-001) to the cell pellet after tapping. Mix well and incubate for 5 min. at room temperature.
- 3) Add 40  $\mu$ L of the primary antibody at the concentration as suggested in the **APPLICATIONS** diluted in the washing buffer. Mix well and incubate for 30 min. at room temperature. (Optimization of antibody concentration or incubation condition is recommended if necessary.)
- 4) Wash the cells 2 times with 1 mL of washing buffer.
- 5) Resuspend the cells with 500  $\mu$ L of the washing buffer and analyze by a flow cytometer.

(Positive control for Flow cytometry; Transfectant)



### ***Flow cytometric detection of mouse GP2 in transfectant***

Cell

Upper: Mouse GP2/293T

Lower: Parental cell (293T)

Antibody

Open: D278-A48

Closed: Isotype control (M081-A48)