

MONOCLONAL ANTIBODY

# Anti-His-tag mAb-Alexa Fluor® 594

Code No.	Clone	Subclass	Quantity	Concentration
D291-A59	OGHis	Mouse IgG2a κ	50 μL	1 mg/mL

**BACKGROUND:** The His-tag (6xHis-tag) is one of the most common tags used to facilitate the purification of recombinant proteins. Metal chelate affinity chromatography is widely used for purification of His-tagged proteins. This specific antibody is useful tool for monitoring of the His-tagged proteins, and recognizes His-tags placed at N-terminal, C-terminal, and internal regions of the recombinant proteins.

**SOURCE:** This antibody was purified from hybridoma (clone OGHIS) supernatant using protein A agarose. This hybridoma was established by fusion of mouse myeloma cell SP-1 with Balb/c mouse splenocyte immunized with 6xHis tagged protein.

**FORMULATION:** 50 μg of IgG in 50 μL volume of PBS containing 1% BSA and 0.1% ProClin150.

**STORAGE:** This antibody solution is stable for one year from the date of purchase when stored at 4°C.

**REACTIVITY:** This antibody recognizes His-tagged protein on Immunocytochemistry.

## APPLICATION:

Immunocytochemistry; 1 μg/mL

\*Please refer to the data sheet (MBL, code no. D291-3) for other applications.

Detailed procedure is provided in the following **PROTOCOL**.

## INTENDED USE:


For Research Use Only. Not for use in diagnostic procedures.

## REFERENCE:

1) Ungerer, C., et al., *Stem Cells Dev.* **23**, 755-766 (2014) [IC]

## RELATED PRODUCTS:

Please visit our website at <https://ruo.mbl.co.jp/>.



### **Immunocytochemical detection of His-tagged calnexin expressed in HeLa using D291-A59.**

Green: Anti-His-tag mAb-Alexa Fluor® 594 (MBL, code no. D291-A59)

Blue: DAPI

The descriptions of the following protocols are examples. Each user should determine the appropriate condition.

## PROTOCOL:

### Immunocytochemistry

- 1) Culture the cells in the appropriate condition on a glass slide. (For example, spread  $1 \times 10^4$  cells of transfectant cells for one slide, then incubate in a CO<sub>2</sub> incubator overnight.)
- 2) Wash the cells 3 times with PBS.
- 3) Fix the cells by immersing the slide in PBS containing 4% paraformaldehyde for 10 minutes at room temperature.
- 4) Wash the glass slide 3 times with PBS.
- 5) Immerse the slide in PBS containing 0.2% Triton X-100 for 10 minutes at room temperature.
- 6) Wash the glass slide 3 times with PBS.
- 7) Add the primary antibody diluted with PBS containing 2% FCS as suggested in the **APPLICATION** onto the cells and incubate for 30 minutes at room temperature. (Optimization of antibody concentration or incubation condition is recommended if necessary.)
- 8) Wash the glass slide twice with PBS.
- 9) Counter stain with DAPI for 5 minutes at room temperature.
- 10) Wash the glass slide twice with PBS.
- 11) Wipe excess buffer off the slide but take care not to touch the cells. Never leave the cells to dry.
- 12) Promptly add mounting medium onto the slide, then put a cover slip on it.

## LABEL LICENSES:

This product is provided under an agreement between LIFE TECHNOLOGIES Corporation, and Medical & Biological Laboratories Co., LTD. Alexa Fluor® is a registered trademark of Molecular Probes, Inc.

210928-5