

Anti-IDH2-R172W (Human) mAb

| | |
|--------------------|---|
| CODE No. | D338-3 |
| CLONALITY | Monoclonal |
| CLONE | WMab-1 |
| ISOTYPE | Rat IgG2b κ |
| QUANTITY | 100 μ L, 1 mg/mL |
| SOURCE | Purified IgG from hybridoma supernatant |
| FORMURATION | PBS containing 50% Glycerol (pH 7.2). No preservative is contained. |
| STORAGE | This antibody solution is stable for one year from the date of purchase when stored at -20°C. |

APPLICATION-CONFIRMED

Western blotting 1 μ g/mL for chemiluminescence detection system

APPLICATIONS-REPORTED

Immunohistochemistry Reference 1)

Immunocytochemistry Reference 1)

SPECIES CROSS REACTIVITY on WB

| Species | Human | Mouse | Rat | Hamster |
|------------|---------------------|------------|------------|------------|
| Sample | Recombinant protein | Not tested | Not tested | Not tested |
| Reactivity | + | | | |

Entrez Gene ID 3418 (Human)

REFERENCES
1) Kato, Y. and Kaneko, M. K., *Biochem Biophys Res Commun.* In press (2013) [WB, IHC, IC]
2) Parsons, D. W., *et al.*, *Science* **321**, 1807-1812 (2008)

For more information, please visit our web site <http://ruo.mbl.co.jp/>


RELATED PRODUCTS

- D338-3 Anti-IDH2-R172W (Human) mAb (WMab-1)
- D311-3 Anti-IDH2 mAb (RMab-22)
- D330-3 Anti-IDH2 mAb (KrMab-3)
- D328-3 Anti-IDH2-R172K (Human) mAb (KMab-1)
- D337-3 Anti-IDH2-R172M (Human) mAb (MMab-1)
- D336-3 Anti-IDH1 (Human) mAb (RcMab-1)
- D309-3 Anti-IDH1 mAb (RMab-3)
- D299-3 Anti-IDH1-R132H (Human) mAb (HMab-1)
- D300-3 Anti-IDH1-R132S (Human) mAb (SMab-1)
- D331-3 Anti-IDH1-R132G (Human) mAb (GMab-r1)

SDS-PAGE & Western blotting

- 1) The recombinant protein is dissolved in Laemmli's sample buffer at 10 µg/mL.
- 2) Boil the samples for 3 min. and centrifuge. Load 10 µL of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel (12.5% acrylamide) for electrophoresis.
- 3) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm² for 1 hr. in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 4) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) for 1 hr. at room temperature.
- 5) Incubate the membrane with primary antibody diluted with 1% skimmed milk (in PBS, pH 7.2) as suggested in the **APPLICATION** for 1 hr. at room temperature. (The concentration of antibody will depend on the conditions.)
- 6) Wash the membrane with PBS-T (0.05% Tween-20 in PBS) (5 min. x 3 times).
- 7) Incubate the membrane with the 1:10,000 anti-IgG (Rat) pAb-HRP (MBL; code no. IM-0825) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hr. at room temperature.
- 8) Wash the membrane with PBS-T (5 min. x 3 times).
- 9) Wipe excess buffer on the membrane, and then incubate it with appropriate chemiluminescence reagent for 1 min. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 10) Expose to an X-ray film in a dark room for 3 min. Develop the film as usual. The condition for exposure and development may vary.

(Positive control for Western blotting; recombinant protein)



Western blot analysis of IDH2-R172W

Lane 1: IDH2 (Wild type)
Lane 2: IDH2-R172K
Lane 3: IDH2-R172W
Lane 4: IDH2-R172M

Immunoblotted with Anti-IDH2-R172W mAb (D338-3)