

Anti-Reelin (Mouse) mAb

CODE No.	D351-3
CLONALITY	Monoclonal
CLONE	2F3
ISOTYPE	Mouse IgG1 κ
QUANTITY	100 μ L, 1 mg/mL
SOURCE	Purified IgG from hybridoma supernatant
IMMUNOGEN	KLH conjugated synthetic peptide, YEKPAFDYC (corresponding to amino acid residues 1241-1248 of mouse Reelin).
REACTIVITY	This antibody does not react with proteolytic fragment of mouse Reelin. Please see the reference 1) for more details.
FORMULATION	PBS containing 50% Glycerol (pH 7.2). No preservative is contained.
STORAGE	This antibody solution is stable for one year from the date of purchase when stored at -20°C.

APPLICATION-CONFIRMED

Western blotting 1 μ g/mL

APPLICATION-REPORTED

Immunohistochemistry Reference 1)

SPECIES CROSS REACTIVITY on WB

Species	Human	Mouse	Rat	Hamster
Sample	Not tested	Culture sup of transfectant	Not tested	Not tested
Reactivity		+		

Entrez Gene ID 19699 (Mouse)

REFERENCE 1) Koie, M., *et al.*, *J. Biol. Chem.* **289**, 12922-12930 (2014) [WB, IHC-fr]

RELATED PRODUCTS


For more information, please visit our web site <https://ruo.mbl.co.jp/>.

The descriptions of the following protocols are examples. Each user should determine the appropriate condition.

SDS-PAGE & Western blotting

- 1) Boil the sample for 2 min. and centrifuge. Load 20 μ L of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel (7.5% acrylamide) for electrophoresis.
- 2) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm² for 1 hr. in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% methanol). See the manufacturer's manual for precise transfer procedure.
- 3) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) overnight at 4°C.
- 4) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 min. x 3).
- 5) Incubate the membrane with primary antibody diluted with 1% skimmed milk (in PBS, pH 7.2) as suggested in the **APPLICATION** for 1 hr. at room temperature. (The concentration of antibody will depend on the conditions.)
- 6) Wash the membrane with PBS-T (5 min. x 3)
- 7) Incubate the membrane with the 1:10,000 of Anti-IgG (H+L chain) (Mouse) pAb-HRP (MBL; code no. 330) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hr. at room temperature.
- 8) Wash the membrane with PBS-T (5 min. x 3).
- 9) Wipe excess buffer on the membrane, and then incubate it with appropriate chemiluminescence reagent for 1 min. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 10) Expose to an X-ray film in a dark room for 5 min. Develop the film as usual. The condition for exposure and development may vary.

(Positive control for Western blotting; Culture sup of transfectant)



Western blotting analysis of mouse Reelin

Sample: Culture sup of NR3/Myc-His transfectant

Immunoblotted with Anti-Reelin (Mouse) mAb (D351-3)

The sample was kindly provided by Dr. Mitsuharu Hattori.
(Department of Biomedical Science, Graduate School of
Pharmaceutical Sciences, Nagoya City University)