

 **My select** sampler set

Anti-GABARAP mAb

| Code No. | Clone | Subclass | Quantity | Concentration |
|----------|-------|------------|----------|---------------|
| M135-3MS | 1F4 | Mouse IgG1 | 20 µL | 1 mg/mL |

BACKGROUND: Autophagy is a process of intracellular bulk degradation in which cytoplasmic components including organelles are sequestered within double-membrane vesicles that deliver the contents to the lysosome/vacuole for degradation. LC3, GABARAP (γ -aminobutyric-acid-type-A-receptor-associated protein), and GATE-16 (Golgi-associated ATPase enhancer of 16 kDa) have been identified as a homologue of yeast Atg8. These homologues have been characterized as modifiers in reactions mediated by hAtg7 (an E1-like enzyme) and hAtg3 (an E2-like enzyme) as in yeast Atg8 lipidation. These homologues also generate form II, which are recovered in membrane fractions. Generation of the form II correlates with autophagosome association. These results suggest that all mammalian Atg8 homologues receive a common modification to associate with autophagosomal membrane as the form II.

SOURCE: This antibody was purified from hybridoma (clone 1F4) supernatant using protein A agarose. This hybridoma was established by fusion of mouse myeloma cell P3U1 with Balb/c mouse lymphocyte immunized with the N-terminus of human GABARAP.

FORMULATION: 20 µg IgG in 20 µL volume of PBS containing 50% glycerol, pH 7.2. No preservative is contained.

STORAGE: This antibody solution is stable for one year from the date of purchase when stored at -20°C.

REACTIVITY: This antibody reacts with GABARAP on Western blotting.

APPLICATIONS:

Western blotting; 1 µg/mL for chemiluminescence detection system

Immunoprecipitation; Not tested

Immunohistochemistry; Not tested

Immunocytochemistry; Not tested

Flow cytometry; Not tested

Detailed procedure is provided in the following **PROTOCOL**.

INTENDED USE:


For Research Use Only. Not for use in diagnostic procedures.

SPECIES CROSS REACTIVITY:

| Species | Human | Mouse | Rat | Hamster |
|------------------|------------------|---------|------------|---------|
| Cells | 293T, HeLa, Raji | NIH/3T3 | Rat1, PC12 | CHO |
| Reactivity on WB | + | + | + | + |

REFERENCES:

- 1) Klionsky, D. J., *et al.*, *J. Cell Sci.* **118**, 7-18 (2005)
- 2) Tanida, I., *et al.*, *J. Biol. Chem.* **277**, 13739-13744 (2002)



Western blot analysis of GABARAP expression in 293T (1), HeLa (2), Raji (3), NIH/3T3 (4), Rat1 (5), PC12 (6) and CHO (7) using M135-3.

The descriptions of the following protocols are examples. Each user should determine the appropriate condition.

PROTOCOL:

SDS-PAGE & Western Blotting

- 1) Wash the 1×10^7 cells 3 times with PBS and suspend with 1 mL of Laemmli's sample buffer.
- 2) Boil the samples for 2 minutes and centrifuge. Load 10 µL of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel for electrophoresis.
- 3) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm² for 1 hour in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 4) To reduce nonspecific binding, place the membrane in 10% skimmed milk (in PBS, pH 7.2) overnight at 4°C.
- 5) Incubate the membrane with primary antibody diluted with PBS, pH 7.2 containing 1% skimmed milk as suggest

in the **APPLICATIONS** for 1 hour at room temperature.
(The concentration of antibody will depend on condition.)

- 6) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 minutes x 3 times).
- 7) Incubate the membrane with the 1:10,000 HRP-conjugated anti-mouse IgG (MBL; code no. 330) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hour at room temperature.
- 8) Wash the membrane with PBS-T (5 minutes x 3 times).
- 9) Wipe excess buffer on the membrane, then incubate it with appropriate chemiluminescence reagent for 1 minute. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 10) Expose to an X-ray film in a dark room for 3 minutes. Develop the film as usual. The condition for exposure and development may vary.

(Positive controls for Western blotting; 293T, HeLa, Raji, NIH/3T3, Rat1, PC12, CHO)

RELATED PRODUCTS:

Please visit our web site <https://ruo.mbl.co.jp/>