

Smart-IP Series

Anti-HA-tag mAb-Magnetic Beads

CODE No.	M180-11
CLONALITY	Monoclonal
CLONE	TANA2
ISOTYPE	Mouse IgG2b κ
QUANTITY	20 tests (Slurry: 1 mL)
SOURCE	Purified IgG from hybridoma supernatant
IMMUNOGEN	KLH conjugated synthetic peptide, YPYDVPDYA (HA-tag)
REACTIVITY	This antibody reacts with N-terminal and C-terminal HA-tagged proteins.
FORMULATION	10 mg magnetic beads in 1 mL PBS/0.1% BSA/0.1% ProClin 150
STORAGE	This beads suspension is stable for one year from the date of purchase when stored at 4°C.

APPLICATION-CONFIRMED

Immunoprecipitation 50 μ L of beads slurry/sample

*The purification capacity of Anti-HA-tag mAb-Magnetic Beads varies depending upon the characteristics of a HA-tagged protein. For example, 50 μ L of beads slurry binds 4.8 μ g of a HA-tagged protein (35 kDa).

APPLICATION-REPORTED

RNP immunoprecipitation Reference 3) and 4)

REFERENCES

- 1) Kato, K., *et al.*, *Nat. Commun.* **9**, 2448 (2018) [IP]
- 2) Shichino, Y., *et al.*, *Elife* **7**, e32155 (2018) [IP]
- 3) Eiken, H. M., *et al.*, *Nat. Commun.* **8**, 1574 (2017) [RIP]
- 4) Jeong, H. W., *et al.*, *Nat. Commun.* **8**, 726 (2017) [RIP]

For more information, please visit our web site <http://ruo.mbl.co.jp/>

RELATED PRODUCTS

Smart-IP series

3190	Magnetic Rack
M180-11	Anti-HA-tag mAb-Magnetic Beads (TANA2)
M132-11	Anti-HA-tag mAb-Magnetic Beads (5D8)
M185-11	Anti-DDDDK-tag mAb-Magnetic Beads (FLA-1)
M047-11	Anti-Myc-tag mAb-Magnetic Beads (PL14)
D291-11	Anti-His-tag mAb-Magnetic Beads (OGHis)
D153-11	Anti-GFP mAb-Magnetic Beads (RQ2)
M165-11	Anti-RFP mAb-Magnetic Beads (3G5)
M167-11	Anti-V5-tag mAb-Magnetic Beads (1H6)
M215-11	Anti-V5-tag mAb-Magnetic Beads (OZA3)
M198-9	Anti-E-tag mAb-Magnetic beads (21D11)
D058-11	Anti-Multi Ubiquitin mAb-Magnetic Beads (FK2)
M075-11	Mouse IgG1 (isotype control)-Magnetic Beads
M076-11	Mouse IgG2a (isotype control)-Magnetic Beads
M077-11	Mouse IgG2b (isotype control)-Magnetic Beads
M081-11	Rat IgG2a (isotype control)-Magnetic Beads
M180-10	Anti-HA-tag mAb-Magnetic Agarose (TANA2)
M132-10	Anti-HA-tag mAb-Magnetic Agarose (5D8)
M185-10	Anti-DDDDK-tag mAb-Magnetic Agarose (FLA-1)
M047-10	Anti-Myc-tag mAb-Magnetic Agarose (PL14)
D291-10	Anti-His-tag mAb-Magnetic Agarose (OGHis)
D153-10	Anti-GFP mAb-Magnetic Agarose (RQ2)
M165-10	Anti-RFP mAb-Magnetic Agarose (3G5)
M167-10	Anti-V5-tag mAb-Magnetic Agarose (1H6)
M198-10	Anti-E-tag mAb-Magnetic Agarose (21D11)

Antibodies

M048-3	Anti-GFP mAb (1E4)
D153-3	Anti-GFP mAb (RQ2)
D153-6	Anti-GFP mAb-Biotin (RQ2)
D153-8	Anti-GFP mAb-Agarose (RQ2)
598	Anti-GFP pAb (polyclonal)
598-7	Anti-GFP pAb-HRP-Direct (polyclonal)
PM073	Anti-Renilla GFP pAb (polyclonal)
M208-3	Anti-RFP mAb Cocktail (1G9, 3G5)
M155-3	Anti-RFP mAb (8D6)
M165-3	Anti-RFP mAb (3G5)
M165-8	Anti-RFP mAb-Agarose (3G5)
M204-3	Anti-RFP mAb (1G9)
M204-7	Anti-RFP mAb-HRP-Direct (1G9)
PM005	Anti-RFP pAb (polyclonal)
PM005-7	Anti-RFP pAb-HRP-Direct (polyclonal)
M180-3	Anti-HA-tag mAb (TANA2) (200 µL)
M180-6	Anti-HA-tag mAb-Biotin (TANA2)
M180-7	Anti-HA-tag mAb-HRP-Direct (TANA2)
561	Anti-HA-tag pAb (polyclonal) (100 µL)
561-7	Anti-HA-tag pAb-HRP-Direct (polyclonal)
561-8	Anti-HA-tag pAb-Agarose (polyclonal)
M132-3	Anti-HA-tag mAb (5D8)
M185-3L	Anti-DDDDK-tag mAb (FLA-1) (1 mL)
M185-7	Anti-DDDDK-tag mAb-HRP-Direct (FLA-1)
PM020	Anti-DDDDK-tag pAb (polyclonal)
PM020-7	Anti-DDDDK-tag pAb-HRP-Direct (polyclonal)
PM020-8	Anti-DDDDK-tag pAb-Agarose (polyclonal)
M192-3	Anti-Myc-tag mAb (My3) (200 µL)
M192-6	Anti-Myc-tag mAb-Biotin (My3)
M047-3	Anti-Myc-tag mAb (PL14)
M047-7	Anti-Myc-tag mAb-HRP-Direct (PL14)

M047-8	Anti-Myc-tag mAb-Agarose (PL14)
562	Anti-Myc-tag pAb (polyclonal) (100 µL)
D291-3	Anti-His-tag mAb (OGHis) (200 µL)
D291-6	Anti-His-tag mAb-Biotin (OGHis)
D291-7	Anti-His-tag mAb-HRP-Direct (OGHis)
D291-8	Anti-His-tag mAb-Agarose (OGHis)
M089-3	Anti-His-tag mAb (6C4)
M136-3	Anti-His-tag mAb (2D8)
PM032	Anti-His-tag pAb (polyclonal)
PM032-8	Anti-His-tag pAb-Agarose (polyclonal)
M167-3	Anti-V5-tag mAb (1H6)
M215-3	Anti-V5-tag mAb (OZA3)
M215-7	Anti-V5-tag mAb-HRP-Direct (OZA3)
PM003	Anti-V5-tag pAb (polyclonal)
PM003-7	Anti-V5-tag pAb-HRP-Direct (polyclonal)
PM003-8	Anti-V5-tag pAb-Agarose (polyclonal)
PM021	Anti-S-tag pAb (polyclonal)
PM070	Anti-E-tag pAb (polyclonal)
PM022	Anti-T7-tag pAb (polyclonal)
563	Anti-VSV-G-tag pAb (polyclonal)
M071-3	Anti-GST-tag mAb (3B2)
M209-3	Anti-GST-tag mAb (GT5)
PM022	Anti-GST-tag pAb (polyclonal)
M095-3	Anti-Luciferase mAb (2D4)
PM016	Anti-Luciferase pAb (polyclonal)
PM047	Anti-Renilla Luciferase pAb (polyclonal)
M094-3	Anti-β-galactosidase mAb (5A3)
PM049	Anti-β-galactosidase pAb (polyclonal)
M091-3	Anti-MBP (Maltose Binding Protein) mAb (1G12)
M013-3	Anti-Thioredoxin (Trx-tag) mAb (2C9)
PM015	Anti-CBD (Chitin Binding Domain) pAb (polyclonal)
PM071	Anti-Calmodulin Binding Protein-tag pAb (polyclonal)
M211-3	Anti-Strep-tag II mAb (4F1)
M214-3	Anti-mini-AID-tag mAb (1E4)


Protein Purification Kits

3320	HA-tagged Protein PURIFICATION KIT
3321	HA-tagged Protein PURIFICATION GEL (1 mL)
3325	DDDDK-tagged Protein PURIFICATION KIT
3326	DDDDK-tagged Protein PURIFICATION GEL (Gel: 1 mL x 1, Peptide: 1 mg x 5)
3328	DDDDK-tagged Protein PURIFICATION GEL (5 mL gel)
3325-205	DDDDK-tag peptide (1 mg x 5)
3326K	DDDDK-tagged Protein PURIFICATION CARTRIDGE (1 mL x 1)
3305	c-Myc-tagged Protein MILD PURIFICATION KIT
3306	c-Myc-tagged Protein MILD PURIFICATION GEL (Gel: 1 mL x 1, Peptide: 1 mg x 1)
3306K	c-Myc-tagged Protein PURIFICATION CARTRIDGE (1 mL x 1)
3310	His-tagged Protein PURIFICATION KIT
3311	His-tagged Protein PURIFICATION GEL (Gel: 1 mL x 1, Peptide: 2 mg x 5)
3317	V5-tagged Protein PURIFICATION KIT Ver.2
3318	V5-tagged Protein PURIFICATION GEL Ver.2 (1 mL)
3315-205	V5-tag peptide (2 mg x 5)

Other related antibodies and kits are also available.
Please visit our website at <http://ruo.mbl.co.jp/>

Immunoprecipitation

- 1) Wash 1×10^7 cells 3 times with PBS and suspend with 1 mL of cold Lysis buffer [50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 1% NP-40] containing appropriate protease inhibitors.
- 2) Centrifuge the tube at $12,000 \times g$ for 5 min. at 4°C and transfer the supernatant to another tube.
- 3) Add magnetic beads as suggested in the **APPLICATION** into 300 μL of the supernatant prepared in step 2). Mix well and incubate with gentle agitation for 1 hr. at 4°C .
- 4) Place the tube on the magnetic rack (MBL; code no. 3190) for a few seconds.
- 5) Remove the supernatant.
- 6) Add 1 mL of cold Wash buffer [50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 0.05% NP-40] and resuspend the magnetic beads.
- 7) Place the tube on the magnetic rack for a few seconds.
- 8) Remove the supernatant.
- 9) Repeat Steps 6)-8) 3 times.
- 10) Resuspend the magnetic beads in 20 μL of Laemmli's sample buffer, boil for 5 min., and place the tube on the magnetic rack for a few seconds.
- 11) Load 20 μL of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel and carry out electrophoresis.
- 12) Visualize the protein bands by CBB staining.



Immunoprecipitation of HA-tagged protein

- Sample: HA-tagged β -galactosidase/293T whole cell lysate
Lane 1: Input (10 μL /lane)
Lane 2: Post-IP beads of Anti-HA-tag mAb (M180-11)