

**For Research Use Only.**  
**Not for use in diagnostic procedures.**



 **My select** sampler set

## Anti-Atg14 (Human) mAb

<b>CODE No.</b>	M184-3MS
<b>CLONALITY</b>	Monoclonal
<b>CLONE</b>	4H8
<b>ISOTYPE</b>	Mouse IgG2a $\kappa$
<b>QUANTITY</b>	20 $\mu$ L, 1 mg/mL
<b>SOURCE</b>	Purified IgG from hybridoma supernatant
<b>IMMUNOGEN</b>	Human Atg14, 167-404 aa (recombinant)
<b>FORMURATION</b>	1 mg/mL in PBS containing 50% Glycerol (pH 7.2). No preservative is contained.
<b>STORAGE</b>	This antibody solution is stable for one year from the date of purchase when stored at -20°C.

### APPLICATIONS-CONFIRMED

<u>Western blotting</u>	1 $\mu$ g/mL for chemiluminescence detection system
<u>Immunoprecipitation</u>	2 $\mu$ g/300 $\mu$ L of cell extract from $3 \times 10^6$ cells

### SPECIES CROSS REACTIVITY on WB

Species	Human	Mouse	Rat	Hamster
Cells	HeLa, 293T, A549, Jurkat	NIH/3T3, MEF	PC12	Not Tested
Reactivity	+	-	-	

**Entrez Gene ID** 22863 (Human)

**REFERENCES**

- 1) Zhong, Y., *et al.*, *Nat. Cell Biol.* **11**, 468 (2009)
- 2) Matsunaga, K., *et al.*, *Nat. Cell Biol.* **11**, 385 (2009)
- 3) Itakura, E., *et al.*, *Mol. Biol. Cell* **19**, 5360 (2008)

For more information, please visit our web site <http://ruo.mbl.co.jp/>



MEDICAL & BIOLOGICAL LABORATORIES CO., LTD.  
URL <http://ruo.mbl.co.jp/>  
e-mail [support@mbi.co.jp](mailto:support@mbi.co.jp), TEL 052-238-1904

**RELATED PRODUCTS**Antibodies

PM036	Anti-LC3 pAb	[WB, IP, IC, IHC, FCM]
M152-3	Anti-LC3 mAb (4E12)	[WB, IP, IC, FCM, EM]
M186-3	Anti-LC3 mAb (8E10)	[WB]
PD014	Anti-LC3 pAb	[WB]
PD015	Anti-LC3 pAb	[IC]
PM046	Anti-LC3 pAb	[WB, IC]
M115-3	Anti-LC3 mAb (51-11)	[WB]
PM045	Anti-p62 (SQSTM1) pAb	
M162-3	Anti-p62 (SQSTM1) (Human) mAb (5F2)	
M162-A48	Anti-p62 (SQSTM1) (Human) mAb -Alexa Fluor®488 (5F2)	
M162-A59	Anti-p62 (SQSTM1) (Human) mAb -Alexa Fluor®594 (5F2)	
M162-A64	Anti-p62 (SQSTM1) (Human) mAb -Alexa Fluor®647 (5F2)	
PM066	Anti-p62 C-terminal pAb	
PD017	Anti-Beclin 1 pAb	
PM037	Anti-GABARAP pAb	
M135-3	Anti-GABARAP mAb (1F4)	
PM038	Anti-GATE-16 pAb	
PD041	Anti-Atg2A pAb	
PM034	Anti-Atg3 pAb	
M133-3	Anti-Atg3 mAb (3E8)	
M134-3	Anti-Atg4B mAb (9H5)	
PM050	Anti-Atg5 pAb	
M153-3	Anti-Atg5 mAb (4D3)	
PM039	Anti-Atg7 (Human) pAb	
PD042	Anti-Atg9A pAb	
M151-3	Anti-Atg10 (Human) mAb (5A7)	
M154-3	Anti-Atg12 (Human) mAb (6E5)	
PD036	Anti-Atg13 (Human) pAb	
M183-3	Anti-Atg13 mAb (5G4)	
PD026	Anti-Atg14 pAb	
M184-3	Anti-Atg14 (Human) mAb (4H8)	
PM040	Anti-Atg16L pAb	
M150-3	Anti-Atg16L mAb (1F12)	
M160-3	Anti-UVRAG mAb (1H4)	
PD027	Anti-Rubicon (Human) pAb	
M170-3	Anti-Rubicon (Human) mAb (1H6)	
PM069	Anti-NRF2 pAb	
M200-3	Anti-NRF2 mAb	
PD037	Anti-Tel2 pAb	
PM072	Anti-VMP1 pAb	

WB: Western blotting  
IP: Immunoprecipitation  
IC: Immunocytochemistry  
IHC: Immunohistochemistry  
FCM: Flow cytometry  
EM: Immuno-electron microscopy

Other related antibodies and kits are also available.  
Please visit our web site at <http://ruo.mbl.co.jp>


Kits

8485	Autophagy Ab Sampler Set
PM036-PN	Positive control for anti-LC3 antibody

**SDS-PAGE & Western blotting**

- 1) Wash  $1 \times 10^7$  cells 3 times with PBS and suspends them in 1 mL of Laemmli's sample buffer, then sonicate briefly (up to 10 seconds).
- 2) Boil the samples for 3 minutes and centrifuge. Load 10  $\mu$ L of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel (12.5% acrylamide) for electrophoresis.
- 3) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm<sup>2</sup> for 1 hour in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 4) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) for overnight at 4°C.
- 5) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 minutes x 3 times).
- 6) Incubate the membrane with primary antibody diluted with PBS, pH 7.2 containing 1% skimmed milk as suggested in the **APPLICATIONS** for 1 hour at room temperature. (The concentration of antibody will depend on the conditions.)
- 7) Wash the membrane with PBS-T (5 minutes x 3 times).
- 8) Incubate the membrane with the 1:10,000 HRP-conjugated anti-mouse IgG (MBL; code no. 330) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hour at room temperature.
- 9) Wash the membrane with PBS-T (5 minutes x 3 times).
- 10) Wipe excess buffer on the membrane, and then incubate it with appropriate chemiluminescence reagent for 1 minute. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 11) Expose to an X-ray film in a dark room for 5 minute. Develop the film as usual. The condition for exposure and development may vary.

(Positive controls for Western blotting; human Atg14 transfectant, HeLa, 293T, A549, and Jurkat)




**Western blot analysis of Atg14**

Lane 1: human Atg14/293T  
Lane 2: HeLa  
Lane 3: 293T  
Lane 4: A549  
Lane 5: Jurkat  
Immunoblotted with M184-3

### **Immunoprecipitation**

- 1) Wash  $1 \times 10^7$  cells 2 times with PBS and resuspend them with 1 mL of ice-cold Lysis buffer (50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 0.05% NP-40) containing appropriate protease inhibitors, then sonicate briefly (up to 20 seconds).
- 2) Centrifuge the tube at 12,000 x g for 5 minutes at 4°C and transfer the supernatant to another tube.
- 3) Mix 20  $\mu$ L of 50% protein A agarose beads slurry resuspended in 300  $\mu$ L of IP buffer (10 mM Tris-HCl (pH 8.0), 500 mM NaCl, 0.1% NP-40) with primary antibody as suggested in the **APPLICATIONS**. Incubate with gently agitation for 1 hour at room temperature.
- 4) Wash the beads 3 times with 1 mL of IP buffer.
- 5) Add 300  $\mu$ L of cell lysate (prepared sample of step 2), then incubate with gentle agitation for 1 hour at room temperature.
- 6) Wash the beads 5 times with 1 mL of Lysis buffer.
- 7) Resuspend the beads in 20  $\mu$ L of Laemmli's sample buffer, boil for 2 minutes and centrifuge.
- 8) Load 20  $\mu$ L of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel (12.5% acrylamide) for electrophoresis.
- 9) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm<sup>2</sup> for 1 hour in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 10) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) for overnight at 4°C.
- 11) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 minutes x 3 times).
- 12) Incubate the membrane with primary antibody diluted with PBS, pH 7.2 containing 1% skimmed milk as suggested in the **APPLICATIONS** for 1 hour at room temperature. (The concentration of antibody will depend on the conditions.)
- 13) Wash the membrane with PBS-T (5 minutes x 3 times).
- 14) Incubate the membrane with the 1:10,000 HRP-conjugated anti-mouse IgG (MBL; code no. 330) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hour at room temperature.
- 15) Wash the membrane with PBS-T (5 minutes x 3 times).
- 16) Wipe excess buffer on the membrane, and then incubate it with appropriate chemiluminescence reagent for 1 minute.
- 17) Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 18) Expose to an X-ray film in a dark room for 5 minutes. Develop the film as usual. The condition for exposure and development may vary.

(Positive controls for Immunoprecipitation; HeLa)



### ***Immunoprecipitation of Atg14 from HeLa***

Lane 1: IP with isotype control (M076-3)  
Lane 2: IP with M184-3  
Immunoblotted with M184-3