

**For Research Use Only.  
Not for use in diagnostic procedures.**



## Anti-Parkin mAb

**CODE No.** M230-3

**CLONALITY** Monoclonal  
**CLONE** Par6  
**ISOTYPE** Mouse IgG2a κ  
**QUANTITY** 100 μL, 1 mg/mL

**SOURCE** Purified IgG from hybridoma supernatant  
**IMMUNOGEN** Human Parkin, full-length (recombinant)  
**FORMURATION** PBS containing 50% Glycerol (pH 7.2). No preservative is contained.  
**STORAGE** This antibody solution is stable for one year from the date of purchase when stored at -20°C.

### APPLICATION-CONFIRMED

Western blotting 1 μg/mL for chemiluminescence detection system

### SPECIES CROSS REACTIVITY on WB

Species	Human*	Mouse	Rat	Hamster
Samples	Transfectant	Brain lysate	Brain lysate, PC12	Not tested
Reactivity	+	+	+	

\*This antibody does not react with HeLa or HEK293T cells.

**Entrez Gene ID** 5071 (Human), 50873 (Mouse), 56816 (Rat)

For more information, please visit our web site <http://ruo.mbl.co.jp/>



## RELATED PRODUCTS

M230-3 Anti-Parkin mAb (Par6)  
M224-3 Anti-KEAP1 mAb (KP1)  
M200-3 Anti-NRF2 mAb (1F2)  
PM069 Anti-NRF2 pAb  
M162-3 Anti-p62 (SQSTM1) (Human) mAb (5F2)  
M162-A48 Anti-p62 (SQSTM1) (Human) mAb  
-Alexa Fluor<sup>®</sup> 488 (5F2)  
M162-A59 Anti-p62 (SQSTM1) (Human) mAb  
-Alexa Fluor<sup>®</sup> 594 (5F2)  
M162-A64 Anti-p62 (SQSTM1) (Human) mAb  
-Alexa Fluor<sup>®</sup> 647 (5F2)  
PM045 Anti-p62 (SQSTM1) pAb  
PM066 Anti-p62 C-terminal pAb  
PM066-7 Anti-p62 C-terminal pAb-HRP-Direct  
M217-3 Anti-Phospho-p62 (SQSTM1) (Ser351) mAb (5D5)  
PM074 Anti-Phospho-p62 (SQSTM1) (Ser351) pAb  
D343-3 Anti-Phospho-p62 (SQSTM1) (Ser403) mAb (4F6)  
D344-3 Anti-Phospho-p62 (SQSTM1) (Ser403) mAb (4C8)  
M152-3 Anti-LC3 mAb (4E12) [WB, IP, IC, FCM, EM]  
M186-3 Anti-LC3 mAb (8E10) [WB]  
M186-7 Anti-LC3 mAb-HRP-Direct (8E10)  
PM036 Anti-LC3 pAb [WB, IP, IC, IHC, FCM]  
PD014 Anti-LC3 pAb [WB]  
PD017 Anti-Becn1 pAb  
PM037 Anti-GABARAP pAb  
M135-3 Anti-GABARAP mAb (1F4)  
PM038 Anti-GATE-16 pAb  
PD041 Anti-Atg2A pAb  
PM034 Anti-Atg3 pAb  
M133-3 Anti-Atg3 mAb (3E8)  
M134-3 Anti-Atg4B mAb (9H5)  
PM050 Anti-Atg5 pAb  
M153-3 Anti-Atg5 mAb (4D3)  
PM039 Anti-Atg7 (Human) pAb  
PD042 Anti-Atg9A pAb  
M151-3 Anti-Atg10 (Human) mAb (5A7)  
M154-3 Anti-Atg12 (Human) mAb (6E5)  
PD036 Anti-Atg13 (Human) pAb  
M183-3 Anti-Atg13 mAb (5G4)  
PD026 Anti-Atg14 pAb  
M184-3 Anti-Atg14 (Human) mAb (4H8)  
PM040 Anti-Atg16L pAb  
M150-3 Anti-Atg16L mAb (1F12)  
M160-3 Anti-UVRAG mAb (1H4)  
PD027 Anti-Rubicon (Human) pAb  
M170-3 Anti-Rubicon (Human) mAb (1H6)  
PD037 Anti-Tel2 pAb  
PM072 Anti-VMP1 pAb  
PM076 Anti-Syntaxin-17 (Human) pAb  
M212-3 Anti-Syntaxin-17 (Human) mAb (2F8)  
  
M175-3 Anti- $\alpha$ -Tubulin mAb (2F9)  
M175-A48 Anti- $\alpha$ -Tubulin mAb-Alexa Fluor<sup>®</sup> 488 (2F9)  
M175-A59 Anti- $\alpha$ -Tubulin mAb-Alexa Fluor<sup>®</sup> 594 (2F9)  
M175-A64 Anti- $\alpha$ -Tubulin mAb-Alexa Fluor<sup>®</sup> 647 (2F9)  
PM054 Anti- $\alpha$ -Tubulin pAb  
PM054-7 Anti- $\alpha$ -Tubulin pAb-HRP-Direct  
M176-3 Anti-EEA1 mAb (3C10)  
M176-A48 Anti-EEA1 mAb-Alexa Fluor<sup>®</sup> 488 (3C10)

M176-A59 Anti-EEA1 mAb-Alexa Fluor<sup>®</sup> 594 (3C10)  
M176-A64 Anti-EEA1 mAb-Alexa Fluor<sup>®</sup> 647 (3C10)  
PM062 Anti-EEA1 pAb  
M178-3 Anti-Calnexin mAb (4F10)  
M178-A48 Anti-Calnexin mAb-Alexa Fluor<sup>®</sup> 488 (4F10)  
M178-A59 Anti-Calnexin mAb-Alexa Fluor<sup>®</sup> 594 (4F10)  
M178-A64 Anti-Calnexin mAb-Alexa Fluor<sup>®</sup> 647 (4F10)  
PM060 Anti-Calnexin pAb  
M181-3 Anti-KDEL mAb (1D5)  
PM059 Anti-KDEL pAb  
M179-3 Anti-GM130 mAb (5G8)  
M179-A48 Anti-GM130 mAb-Alexa Fluor<sup>®</sup> 488 (5G8)  
M179-A59 Anti-GM130 mAb-Alexa Fluor<sup>®</sup> 594 (5G8)  
M179-A64 Anti-GM130 mAb-Alexa Fluor<sup>®</sup> 647 (5G8)  
PM061 Anti-GM130 pAb  
PM063 Anti-COX4 pAb  
PM064 Anti-Lamin B1 pAb

### Kits

8485 Autophagy Ab Sampler Set  
8486 Autophagy Watch  
PM036-PN Positive control for anti-LC3 antibody


WB: Western blotting  
IP: Immunoprecipitation  
IC: Immunocytochemistry  
IHC: Immunohistochemistry  
FCM: Flow cytometry  
EM: Immuno-electron microscopy

Other related antibodies and kits are also available.  
Please visit our website at <http://ruo.mbl.co.jp/>

### **SDS-PAGE & Western blotting**

- 1) Wash  $1 \times 10^7$  cells 3 times with PBS and suspend them in 1 mL of Laemmli's sample buffer, then sonicate briefly (up to 20 sec.).
- 2) Boil the sample for 5 min. and centrifuge.
- 3) Load 10  $\mu$ L of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel (12.5% acrylamide) for electrophoresis.
- 4) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm<sup>2</sup> for 1 hr. in a dry transfer system.
- 5) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) overnight at 4°C.
- 6) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 min. x 3 times).
- 7) Incubate the membrane with primary antibody diluted with 1% skimmed milk (in PBS, pH 7.2) as suggested in the **APPLICATION** for 1 hr. at room temperature. (The concentration of antibody will depend on the conditions.)
- 8) Wash the membrane with PBS-T (5 min. x 3 times).
- 9) Incubate the membrane with 1:10,000 of Anti-IgG (Mouse) pAb-HRP (MBL; code no. 330) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hr. at room temperature.
- 10) Wash the membrane with PBS-T (5 min. x 3 times).
- 11) Wipe excess buffer on the membrane, then incubate it with appropriate chemiluminescence reagent for 1 min. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 12) Expose to an X-ray film in a dark room for 3 min. Develop the film as usual. The condition for exposure and development may vary.

(Positive controls for Western blotting; Rat brain, mouse brain, PC12 and transfectant)



#### ***Western blot analysis of Parkin***

- Lane 1: Rat brain lysate, 20  $\mu$ g
- Lane 2: Mouse brain lysate, 20  $\mu$ g
- Lane 3: PC12
- Lane 4: HeLa
- Lane 5: HEK293T
- Lane 6: Human Parkin/HEK293T

Immunoblotted with Anti-Parkin mAb (M230-3)