

 **My select** sampler set

# Anti-GABARAP pAb

<b>Code No.</b>	<b>Quantity</b>	<b>Form</b>
PM037MS	20 $\mu$ L	Affinity Purified

**BACKGROUND:** Autophagy is a process of intracellular bulk degradation in which cytoplasmic components including organelles are sequestered within double-membrane vesicles that deliver the contents to the lysosome/vacuole for degradation. LC3, GABARAP ( $\gamma$ -aminobutyric-acid-type-A-receptor-associated protein), and GATE-16 (Golgi-associated ATPase enhancer of 16 kDa) have been identified as a homologue of yeast Atg8. These homologues have been characterized as modifiers in reactions mediated by hAtg7 (an E1-like enzyme) and hAtg3 (an E2-like enzyme) as in yeast Atg8 lipidation. These homologues also generate form II, which are recovered in membrane fractions. Generation of the form II correlates with autophagosome association. These results suggest that all mammalian Atg8 homologues receive a common modification to associate with autophagosomal membrane as the form II.

**SOURCE:** This antibody was purified from rabbit serum using affinity column. The rabbit was immunized with carrier protein conjugated synthetic peptide at the N-terminus region of GABARAP.

**FORMULATION:** 20  $\mu$ L volume of PBS containing 50% glycerol, pH 7.2. No preservative is contained.

**STORAGE:** This antibody solution is stable for one year from the date of purchase when stored at  $-20^{\circ}\text{C}$ .

**REACTIVITY:** This antibody reacts with GABARAP on Western blotting and Immunocytochemistry. It does not react with LC3 and GATE16.

## APPLICATIONS:

Western blotting; 1:1,000  
Immunoprecipitation; Not tested  
Immunohistochemistry; Not tested  
Immunocytochemistry; 1:100  
Flow cytometry; Not tested

Detailed procedure is provided in the following **PROTOCOLS**.

## INTENDED USE:

For Research Use Only. Not for use in diagnostic procedures.


## SPECIES CROSS REACTIVITY:

Species	Human	Mouse	Rat	Hamster
Cells	293T, HeLa, Raji, HL-60, Jurkat	NIH/3T3, WR19L	Rat1, PC12	CHO
Reactivity on WB	+	+	+	+


## REFERENCES:

- 1) Klionsky, D. J., *et al.*, *J. Cell Sci.* **118**, 7-18 (2005)
- 2) Tanida, I., *et al.*, *J. Biol. Chem.* **277**, 13739-13744 (2002)

kDa 1 2 3 4 5 6 7 8 9 10



**Western blotting analysis of GABARAP expression in 293T (1), HeLa (2), Raji (3), HL-60 (4), Jurkat (5), NIH/3T3 (6), WR19L (7), Rat1 (8), PC12 (9) and CHO (10) using PM037.**



**Western blotting analysis of GABARAP expression on transfectant (TF) of human Atg8 homologs using PM037.**

## PROTOCOLS


### SDS-PAGE & Western blotting

- 1) Wash the  $1 \times 10^7$  cells 3 times with PBS and suspend with 1 mL of Laemmli's sample buffer.
- 2) Boil the samples for 2 minutes and centrifuge. Load 10  $\mu$ L of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel for electrophoresis.
- 3) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm<sup>2</sup> for 1 hour in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% methanol). See the manufacture's manual for precise transfer procedure.
- 4) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) for overnight at 4°C.
- 5) Incubate the membrane with primary antibody diluted with 1% skimmed milk (in PBS, pH 7.2) as suggested in the **APPLICATIONS** for 1 hour at room temperature. (The concentration of antibody will depend on the conditions.)
- 6) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 minutes x 3).
- 7) Incubate the membrane with the 1:10,000 HRP-conjugated anti-rabbit IgG (MBL, code no. 458) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hour at room temperature.
- 8) Wash the membrane with PBS-T (5 minutes x 3).
- 9) Wipe excess buffer on the membrane, then incubate it with appropriate chemiluminescence reagent for 1 minute.
- 10) Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 11) Expose to an X-ray film in a dark room for 3 minutes.
- 12) Develop the film as usual. The condition for exposure and development may vary.

(Positive controls for Western blotting; 293T, HeLa, Raji, HL60, Jurkat, NIH/3T3, WR19L, Rat1, PC12, CHO)

- 4) Fix the cells by immersing the slide in 4% paraformaldehyde PBS for 10 minutes at room temperature (20~25°C).
- 5) Prepare a wash container such as a 500 mL beaker with a magnetic stirrer. Then wash the fixed cells on the glass slide by soaking the slide with a plenty of PBS in the wash container for 5 minutes. Take care not to touch the cells. Repeat another wash once more.
- 6) Immerse the slide in 100  $\mu$ g/mL of Digitonin for 15 minutes at room temperature.
- 7) Wash the slide in a plenty of PBS as in the step 5).
- 8) Add the primary antibody diluted with PBS as suggested in the **APPLICATIONS** onto the cells and incubate for 1 hour at room temperature. (Optimization of antibody concentration or incubation condition is recommended if necessary.)
- 9) Wash the slide in a plenty of PBS as in the step 5).
- 10) Add 200  $\mu$ L of 1:100 FITC conjugated anti-rabbit IgG (MBL, code no. IM-0833) diluted with PBS onto the cells. Incubate for 30 minutes at room temperature. Keep out light by aluminum foil.
- 11) Wash the slide in a plenty of PBS as in the step 5).
- 12) Wipe excess liquid from slide but take care not to touch the cells. Never leave the cells to dry.
- 13) Promptly add mounting medium onto the slide, then put a cover slip on it.

(Positive control for Immunocytochemistry; NRK)



**Immunocytochemical detection of GABARAP on 4% PFA fixed nutrient normal rat kidney cell line (NRK, left) and starved NRK (right) with PM037.**

### Immunocytochemistry

- 1) Spread the cells in the nutrient condition on a glass slide, then incubate in a CO<sub>2</sub> incubator for one night.
- 2) Remove the culture supernatant by careful aspiration.
- 3) To obtain serum-starved conditions, culture the cells with Hank's solution or DMEM for 2-4 hours at 37°C.