

POLYCLONAL ANTIBODY

cis-Golgi Marker

Anti-GM130 pAb

Code No.
PM061

Quantity
100 μ L

Form
Affinity Purified

BACKGROUND: The Golgi apparatus is a eukaryotic organelle, which is mainly devoted to processing the proteins synthesized in the endoplasmic reticulum (ER). GM130 is a member of the golgin family of coiled-coil proteins that localizes predominantly to the cis-Golgi. GM130 might participate in ER-Golgi traffic.

SOURCE: This antibody was purified from rabbit serum using affinity column. The rabbit was immunized with the synthetic peptide corresponding to C-terminus of GM130.

FORMULATION: 100 μ L volume of PBS containing 50% glycerol, pH 7.2. No preservative is contained.

STORAGE: This antibody solution is stable for one year from the date of purchase when stored at -20°C .

REACTIVITY: This antibody reacts with human GM130 for Western blotting, Immunoprecipitation and Immunocytochemistry.

APPLICATIONS:

Western blotting; 1:1,000

Immunoprecipitation; 2 μ L/300 μ L of cell extract from 1×10^7 cells

Immunohistochemistry; Not tested

Immunocytochemistry; 1:500

Flow cytometry; Not tested

Detailed procedures are provided in the following **PROTOCOLS**.

SPECIES CROSS REACTIVITY:

Species	Human	Mouse*	Rat
Cells	HeLa, 293T, A549	NIH/3T3	Not Tested
Reactivity on WB	+	-	

*This antibody does not react with mouse in Western blotting, but can be used in Immunocytochemistry.


INTENDED USE:

For Research Use Only. Not for use in diagnostic procedures.

REFERENCES:

- 1) Seto, S., *et al.* *PLoS One* **8**, e83324 (2013) [IC]
- 2) Tamaki, H., *et al.*, *FEBS Lett.* **586**, 3064-3070 (2012) [IC]
- 3) Diao, A., *et al.*, *J. Biol. Chem.* **283**, 6957-6967 (2008)
- 4) Alvarez, C., *et al.*, *J. Biol. Chem.* **276**, 2693-2700 (2001)

The descriptions of the following protocols are examples. Each user should determine the appropriate condition.



Western blot analysis of GM130 expression on HeLa (1), 293T (2) and A549 (3) using PM061.


PROTOCOLS:

SDS-PAGE & Western Blotting

- 1) Wash cells (approximately 1×10^7 cells) 3 times with PBS and resuspend them in 1 mL of Laemmli's sample buffer.
- 2) Boil the samples for 2 minutes and centrifuge. Load 10 μ L of sample per lane on a 1-mm-thick SDS-polyacrylamide gel and carry out electrophoresis.
- 3) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm² for 1 hour in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 4) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) for 1 hour at room temperature, or overnight at 4°C.
- 5) Incubate the membrane for 1 hour at room temperature with primary antibody diluted with 1% skimmed milk (in PBS, pH 7.2) as suggested in the **APPLICATIONS**. (The concentration of antibody will depend on the conditions.)
- 6) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 minutes x 3 times).
- 7) Incubate the membrane with 1:10,000 of Anti-IgG (Rabbit) pAb-HRP (MBL; code no. 458) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hour at room temperature.

- 8) Wash the membrane with PBS-T (5 minutes x 3 times).
- 9) Wipe excess buffer off the membrane, and incubate membrane with an appropriate chemiluminescence reagent for 1 minute.
- 10) Remove extra reagent from the membrane by dabbing with a paper towel, and seal it in plastic wrap.
- 11) Expose the membrane onto an X-ray film in a dark room for 3 minutes. Develop the film under usual settings. The conditions for exposure and development may vary.

(Positive controls for Western blotting; HeLa, 293T, A549 and Jurkat)



Immunoprecipitation of GM130 from HeLa with normal rabbit IgG (1) or PM061 (2). After immunoprecipitated with the antibody, immunocomplex was resolved on SDS-PAGE and immunoblotted with PM061.

Immunoprecipitation

- 1) Wash cells (approximately 1×10^7 cells) 3 times with PBS and resuspend them in 1 mL of cold Lysis buffer [50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 0.05% NP-40] containing protease inhibitors at appropriate concentrations. Incubate it at 4°C with rotating for 30 minutes; thereafter, briefly sonicate the mixture (up to 10 seconds).
- 2) Centrifuge the tube at 12,000 x g for 10 minutes at 4°C and transfer the supernatant to another tube.
- 3) Add primary antibody as suggested in the **APPLICATIONS** into 300 μ L of the supernatant. Mix well and incubate with gentle agitation for 30-120 minutes at 4°C. Add 20 μ L of 50% protein A agarose beads resuspended in the cold Lysis buffer. Mix well and incubate with gentle agitation for 60 minutes at 4°C.
- 4) Wash the beads 3-5 times with the cold Lysis buffer (centrifuge the tube at 2,500 x g for 10 seconds).
- 5) Resuspend the beads in 20 μ L of Laemmli's sample buffer, boil for 3-5 minutes, and centrifuge for 5 minutes. Use 10 μ L/lane for the SDS-PAGE analysis.



(See **SDS-PAGE & Western blotting**.)

(Positive control for Immunoprecipitation; HeLa)

Immunocytochemistry

- 1) Culture the cells in the appropriate condition on a glass slide. (for example, spread 1×10^4 cells for one slide, then incubate in a CO₂ incubator for one night.)
- 2) Wash the glass slide 2 times with PBS.
- 3) Fix the cells by immersing the slide in PBS containing 4% paraformaldehyde for 10 minutes at room temperature.
- 4) Wash the glass slide 3 times with PBS.
- 5) Immerse the slide in PBS containing 0.2% Triton X-100 for 10 minutes at room temperature.
- 6) Wash the glass slide 2 times with PBS.
- 7) Add the primary antibody diluted with 2% FCS/PBS as suggested in the **APPLICATIONS** onto the cells and incubate for 1 hour at room temperature (Optimization of antibody concentration or incubation condition are recommended if necessary).
- 8) Wash the glass slide 2 times with PBS.
- 9) Add 100 μ L of 1:500 Alexa Fluor® 488 conjugated anti-rabbit IgG (Invitrogen; code no. A110374) diluted with PBS onto the cells. Incubate for 30 minutes at room temperature. Keep out light by aluminum foil.
- 10) Wash the glass slide 3 times with PBS.
- 11) Counter stain with DAPI for 5 minutes at room temperature.
- 12) Wash the glass slide 2 times with PBS.
- 13) Wipe excess liquid off the slide but take care not to touch the cells. Never leave the cells to dry.
- 14) Promptly add mounting medium onto the slide, then put a cover slip on it.

(Positive control for Immunocytochemistry; A549 and NIH/3T3)



Immunocytochemical detection of GM130 in A549 (upper) and in NIH/3T3 (lower) with PM061.

Green: Anti-GM130 pAb (PM061)
Blue: DAPI counter stain

RELATED PRODUCTS:

Please visit our website at <https://ruo.mbl.co.jp/>.