

**For Research Use Only.**  
**Not for use in diagnostic procedures.**



# Anti-5-hydroxymethylcytosine (5hmC) pAb

<b>CODE No.</b>	PM077
<b>CLONALITY</b>	Polyclonal
<b>ISOTYPE</b>	Rabbit Ig, affinity purified
<b>QUANTITY</b>	100 µL
<b>SOURCE</b>	Purified IgG from rabbit serum
<b>IMMUNOGEN</b>	BSA-conjugated 5-hydroxymethylcytidine
<b>FORMURATION</b>	PBS containing 50% Glycerol (pH 7.2). No preservative is contained.
<b>STORAGE</b>	This antibody solution is stable for one year from the date of purchase when stored at -20°C.

## APPLICATIONS-CONFIRMED

<u>Dot blotting</u>	1:5,000 for chemiluminescence detection system
<u>Hydroxymethylated DNA immunoprecipitation (hMeDIP)</u>	2 µL/sample

For more information, please visit our web site <http://ruo.mbl.co.jp/>



## **RELATED PRODUCTS**

### Antibodies

PM077	Anti-5-hydroxymethylcytosine (5hmC) pAb
MI-11-3	Anti-Bromodeoxyuridine mAb (2B1)
D209-3	Anti-Histone H1 mAb (C14093)
D210-3	Anti-Histone H2A mAb (C10037)
D212-3	Anti-Histone H2B mAb (C14264)
PM006	Anti-Phospho-Histone H3 (Ser28) (Human) pAb
PM006-A48	Anti-Phospho-Histone H3 (Ser28) (Human) pAb -Alexa Fluor <sup>®</sup> 488
CY-M1029	Anti-Acetylated Histone/p53 (Lys382) mAb (TM-5C5)
CY-P1011	Anti-HDAC1 (Histone Deacetylase 1) pAb
CY-P1012	Anti-HDAC2 (Histone Deacetylase 2) pAb
CY-P1015	Anti-Phospho-Histone-H2A.X (Ser139) pAb
PM035	Normal Rabbit IgG

### Kits

5350	MethylHunter 5hmC detection kit
5270-100	MethylHunter MBD1-based Methylated DNA Enrichment Kit
5275-100	MethylHunter MBD1-based Methylated DNA Enrichment Kit 2


Other related antibodies and kits are also available.

Please visit our website at <http://ruo.mbl.co.jp/>

## **Dot blotting**

- 1) Sample preparation:
  - a) Prepare DNA samples by appropriate method (e.g., 5hmC-containing DNA by performing PCR).
  - b) Add 0.1 volumes of 1 M NaOH to the DNA samples.
  - c) Heat the DNA samples at 99°C for 5 min., then quench at 0°C for 5 min.
  - d) Add 0.1 volumes of 6.6 M NH<sub>4</sub>OAc to the DNA samples.
- 2) Blot 1 µL of different concentrations of DNA samples onto a nitrocellulose membrane.
- 3) Cross-link the DNA samples using UV illuminator for 5 min.
- 4) To reduce nonspecific binding, soak the membrane in 5% skimmed milk (in PBS, pH 7.2) for 1 hr. at room temperature.
- 5) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 min. x 3 times).
- 6) Incubate the membrane with primary antibody diluted with 1% skimmed milk (in PBS, pH 7.2) as suggested in the **APPLICATIONS** for 1 hr. at room temperature. (The concentration of antibody will depend on the conditions.)
- 7) Wash the membrane with PBS-T (5 min. x 3 times).
- 8) Incubate the membrane with the 1:5,000 Anti-IgG (Rabbit) pAb-HRP (MBL; code no. 458) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hr. at room temperature.
- 9) Wash the membrane with PBS-T (5 min. x 3 times)
- 10) Wipe excess buffer on the membrane, then incubate it with appropriate chemiluminescence reagent for 1 min. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 11) Expose to an X-ray film in a dark room for 5 min. Develop the film as usual. The condition for exposure and development may vary.

(Positive control for Dot blotting; PCR synthesized DNA containing 5hmC)



### ***Dot blot analysis of 5hmC-containing DNA***

- Lane 1: PCR synthesized DNA containing 114 C residues (414 bp)
- Lane 2: PCR synthesized DNA containing 114 5mC residues (414 bp)
- Lane 3: PCR synthesized DNA containing 114 5hmC residues (414 bp)


Immunoblotted with Anti-5-hydroxymethylcytosine (5hmC) pAb (PM077)

### Hydroxymethylated DNA Immunoprecipitation (hMeDIP)

1) Sample preparation:

- a) Prepare 1 µg of genomic DNA fragment by appropriate method.
  - b) Add 50 pg of PCR synthesized C-, 5mC- or 5hmC-containing DNA as a spike-in control.
  - c) Dissolve the DNA sample in TE buffer [10 mM Tris-HCl (pH 7.4), 1 mM EDTA] and adjust the volume to 20 µL.
  - d) Heat the DNA sample at 99°C for 10 min., then quench at 0°C for 10 min.
- 2) Add 430 µL of TE buffer, 50 µL of 10 x IP buffer [100 mM Na-Phosphate (pH 7.0), 1.4 M NaCl, 0.5 % Triton X-100] and Anti-5-hydroxymethylcytosine (5hmC) pAb (PM077) as suggested in the **APPLICATIONS** or 2 µg of Normal Rabbit IgG (MBL; code no. PM035). Incubate with gentle agitation for 2 hr. at 4°C.
  - 3) During step 2), wash 40 µL of Dynabeads® M-280 Sheep anti-Rabbit IgG (Life Technologies; code no. 11203D) with 800 µL of PBS and place the tube on the magnetic rack (MBL; Code no. 3190) for a few seconds. Discard the supernatant carefully. Resuspend the beads with 50 µL of 1 x IP buffer [10 mM Na-Phosphate (pH 7.0), 140 mM NaCl, 0.05 % Triton X-100].
  - 4) Add 50 µL of washed beads suspension (prepared in step 3)) to DNA and antibody mixture. Incubate with gentle agitation for 2 hr. at 4°C.
  - 5) Place the tube on the magnetic rack for a few seconds and discard the supernatant carefully.
  - 6) For washing the beads, add 700 µL of 1 x IP buffer and incubate the tube with gentle rotation for 10 min. at room temperature.
  - 7) Place the tube on the magnetic rack for a few seconds and discard the supernatant carefully.
  - 8) Repeat 3 times steps 6)-7).
  - 9) Add 250 µL of Proteinase K digestion buffer [50 mM Tris (pH 8.0), 10 mM EDTA, 0.5 % SDS]. Incubate for 1 hr. at 50°C with inversion every 10 min.
  - 10) Perform DNA extraction by Phenol/Chloroform extraction followed by ethanol precipitation.
  - 11) Dissolve the pellet in 60 µL of nuclease-free water.

(Positive control for Dot blotting; PCR synthesized DNA containing 5hmC)



### ***Analysis of immunoprecipitated DNA with Real-time PCR using spike-in control specific primers***

C: PCR synthesized DNA (13 CpG/136 bp)

5mC: PCR synthesized DNA (13 5mCpG/136 bp)

5hmC: PCR synthesized DNA (13 5hmCpG/136 bp)