

For Research Use Only.
Not for use in diagnostic procedures.

RiboCluster Profiler™

Anti-7-methylguanosine (m⁷G) mAb

CODE No.	RN017M
CLONALITY	Monoclonal
CLONE	4141-13
ISOTYPE	Mouse IgG2a κ
QUANTITY	200 μL, 1 mg/mL
SOURCE	Purified IgG from hybridoma supernatant
IMMUNOGEN	Carrier protein-conjugated 7-methylguanosine (m ⁷ G)-Cap analogue
REACTIVITY	This clone reacts with both 5'-terminal and internal 7-methylguanosine (m ⁷ G) in RNA.
FORMULATION	PBS containing 50% Glycerol (pH 7.2). No preservative is contained.
STORAGE	This antibody solution is stable for one year from the date of purchase when stored at -20°C.

APPLICATIONS-CONFIRMED

<u>Dot blotting</u>	1 μg/mL
<u>RNA immunoprecipitation</u>	10 μg/sample
<u>Immunocytochemistry</u>	Can be used.
<u>RNA EISA</u>	Can be used.

APPLICATION-REPORTED

<u>Northwestern blotting</u>	Reference 1)
------------------------------	--------------

REFERENCES


- 1) Lin, S., *et al.*, *Mol. Cell* **71**, 244-255.e5 (2018) [RNA-IP, Northwestern blotting]
- 2) Zorbas, C., *et al.*, *Mol. Biol. Cell*. **26**, 2080-2095 (2015)
- 3) Filonava, L., *et al.*, *Genome Biol.* **16**, 216 (2015)
- 4) Chen, P., *et al.*, *BMC Plant Biol.* **10**, 201 (2010)
- 5) Cubrilo, S., *et al.*, *RNA* **15**, 1492-1497 (2009)
- 6) Cowling, V. H., *Biochem. J.* **425**, 295-302 (2009)
- 7) Maden, B. E. and Salim, M., *J. Mol. Biol.* **88**, 133-152 (1974)

For more information, please visit our website <https://ruo.mbl.co.jp>.

Dot blotting

Dot blotting was performed using DIG Wash and Block Buffer Set (Sigma-Aldrich; code no. 11585762001). For more information, please contact Sigma-Aldrich Co. LLC.

- 1) Sample preparation:
 - a) Prepare RNA samples by appropriate method (e.g., m⁷G-capped RNA by *in vitro* transcription).
 - b) Heat the RNA samples at 80°C for 2 min., then quench at 4°C for 5 min.
- 2) Blot 1 µL of different concentrations of the RNA samples onto a nitrocellulose membrane.
- 3) Cross-link the RNA samples using UV illuminator.
- 4) To reduce nonspecific binding, soak the membrane in Blocking Solution for 30 min. at room temperature.
- 5) Incubate the membrane with primary antibody diluted with Blocking Solution as suggested in the **APPLICATIONS** for 1 hr. at room temperature. (The concentration of antibody will depend on the conditions.)
- 6) Wash the membrane with Washing Buffer (15 min. x 2).
- 7) Incubate the membrane with 1:5,000 of Anti-IgG (Mouse) pAb-HRP (MBL; code no. 330) diluted with Blocking Solution for 1 hr. at room temperature.
- 8) Wash the membrane with Washing Buffer (15 min. x 2).
- 9) Wash the membrane with Washing Buffer (3 min. x 1).
- 10) Wipe excess buffer on the membrane, then incubate it with appropriate chemiluminescence reagent for 1 min. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 11) Expose for 40 sec. with ImageQuant LAS 4000 imaging system (Fujifilm). The condition for exposure and development may vary.



Dot blot analysis of m⁷G-capped RNA

Sample: *In vitro* transcribed RNA from full-length of RN7SK RNA (RefSeq ID: NR_001445)

Immunoblotted with Anti-7-methylguanosine (m⁷G) mAb (MBL; code no. RN017M)

RNA immunoprecipitation

Some buffers and reagents are included in the RIP-Assay Kit *for microRNA* (MBL; code. RN1005). Please also refer to the protocol packaged in the RIP-Assay Kit *for microRNA*.


[Material Preparation]

1. **RNA-IP Buffer (+)** [mi-Lysis Buffer (component of RN1005) containing 1.5 mM DTT and RNase inhibitor]
Before using RNA-IP Buffer (+), RNase inhibitor and DTT are added to mi-Lysis Buffer at the appropriate concentration.
2. **Wash Buffer** [mi-Wash Buffer (component of RN1005) containing 1.5 mM DTT]
Before using Wash Buffer, DTT is added to mi-Wash Buffer at the appropriate concentration.
3. Antibody conjugated Protein G beads
 - A) Mix 20 μ L of 50% protein G agarose beads slurry resuspended in nuclease-free PBS with 600 μ L of mi-Wash Buffer (component of RN1005), and then add Mouse IgG2a (isotype control) (MBL; code no. M076-3) or Anti-7-methylguanosine (m^7G) mAb (RN017M) at the concentration suggested in the **APPLICATIONS**. Incubate with gentle agitation overnight at 4°C.
 - B) Wash the beads 1 time with mi-Lysis Buffer (component of RN1005) containing 1.5 mM DTT.
 - C) Carefully discard the supernatant using a pipettor without disturbing the beads and incubate at 4°C until just before use.
4. Input total RNA
Prepare total RNA samples by appropriate isolation method. Heat-denature the total RNA samples at 80°C for 2 min., then quench at 4°C for more than 5 min.

[Protocol (RNA isolation; 2-step method in RN1005)]



- 1) Add 40 μ g of input total RNA and 500 μ L of RNA-IP Buffer into the tube containing antibody conjugated beads, then incubate with gentle agitation for 3 hr. at 4°C.
- 2) Wash the beads 4 times with 1 mL of Wash Buffer (centrifuge the tube at 2,000 x g for 1 min.).
- 3) Add 250 μ L of Master mix solution (mi-Solution I: mi-Solution II = 10 μ L: 240 μ L). Vortex thoroughly, then spin-down.
- 4) Add 150 μ L of mi-Solution III. Vortex thoroughly.
- 5) Centrifuge the tube at 2,000 x g for 2 min.
- 6) Transfer the supernatant to the new tube containing 2 μ L of mi-Solution IV.
- 7) Add 400 μ L of ice-cold 100% ethanol. Vortex thoroughly, then spin-down. Place at -20°C for 20 min. Centrifuge the tube at 12,000 x g for 10 min. at 4°C, then add 2 μ L of mi-Solution IV to the supernatant in the same tube.
- 8) Add 400 μ L of ice-cold 100% ethanol. Vortex thoroughly, then spin-down. Place at -20°C for 20 min. Centrifuge the tube at 12,000 x g for 10 min. at 4°C.
- 9) Wash the pellet 2 times with 500 μ L of ice-cold 70% ethanol and dry up the pellet for 5-15 min.
- 10) Dissolve the pellets in 20 μ L of nuclease-free water. Quantify the isolated RNA using NanoDrop (Thermo Fisher Scientific Inc.) and check the quality of RNA with Experion (Bio-Rad).

(Positive control for RNA immunoprecipitation; HEK293T total RNA)



RNA immunoprecipitation from *in vitro* transcribed RNA

Sample: 2 μ g of *in vitro* transcribed RNA
from full-length of RN7SK RNA
(RefSeq ID: NR_001445)



(B)

Average of the RNA Quantity (n=2)	
Antibody	RNA (ng)
Anti-7-methylguanosine (m ⁷ G) mAb	921.0
Mouse IgG2a	42.2

RNA immunoprecipitation from HEK293T total RNA

- (A) Characterization of isolated RNA with Experion
- (B) Quantification of isolated RNA with NanoDrop